


Sub

[illegible]

substituents selected from the group consisting of cyano groups, halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a biphenyl group; an α-cyanobenzyl group; an α-cyanobenzyl group substituted with 1 to 5 halogen atoms; a benzyl group substituted with a bicyclo[2.2.1]-hept-5-en-2,3-dicarboxyimidyl group; a benzoyl group; a styryl group; a styryl group substituted with 1 to 5 substituents selected from the group consisting of C₁₋₆ alkoxy groups and di(C₁₋₆ alkyl)amino alkyl groups; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyrimidinyl group; a pyrimidinyl group substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups and C₁₋₆ alkoxy groups; a phthalimidoyl group; a phthalimidoyl group substituted with 1 to 3 halogen atoms; an N-carbazolyl group; a dioxopiperidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups; a phenylsulfonylamino group; a phenylsulfonylamino group substituted with 1 to 3 C₁₋₆ alkyl groups; a C₁₋₆ alkylaminosulfonyl C₁₋₆ alkyl group; a thiadiazolyl group; an oxadiazolyl group; an oxadiazolyl group substituted with a substituted phenyl group wherein the substituents in the substituted phenyl group are 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a pyrrolidinyl group; a pyrazolyl group; a pyrazolyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and trifluoromethyl groups; a furyl group; a furyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₂₋₆ alkoxy carbonyl groups; a thienopyrimidinylthio group; a thienopyrimidinylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a thienopyridylthio group; a thienopyridylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a benzothiazolylthio group; a benzothiazolylthio group substituted with 1 to 3 halogen atoms; a group represented by the formula: -Y-(CR⁶¹R⁶²)_m-(CR⁶³R⁶⁴)_n-R⁷ [wherein Y represents an oxygen or sulfur atom; R⁶¹, R⁶², R⁶³, and R⁶⁴ are identical or different and represent a hydrogen atom, a halogen atom, a C₁₋₄ alkyl group, or a

[illegible]

Sub B¹

alkyl) amino group; a C₂₋₆ alkoxy carbonyl group; or a di(C₁₋₆ alkyl) amino C₁₋₆ alkoxy group; m is an integer of 1 to 6; and n is an integer of 0 to 6]; or a group represented by the formula: -SO₂NR⁸R⁹ [wherein R⁸ and R⁹ are identical or different and represent a hydrogen atom, a C₁₋₁₀ alkyl group, a C₂₋₆ alkanoyl group, an isoxazolyl group, an isoxazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiadiazolyl group, a thiadiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiazolyl group, a thiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyridyl group, a pyridyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyrimidinyl group, a pyrimidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyrimidinyl group substituted with 1 to 3 C₁₋₆ alkoxy groups, a pyridazinyl group, a pyridazinyl group substituted with 1 to 3 C₁₋₆ alkoxy groups, an indazolyl group, or a carbamoyl group mono- or di-substituted with C₁₋₆ alkyl groups, or alternatively, taken together with the nitrogen atom to which they are bonded, form a 3,5-dioxopiperadino group, a pyrrolidinyl group, a piperidino group, or a morpholino group], or alternatively,

the two groups adjacent to each other of R¹ to R⁵, taken together with the benzene ring to which they are bonded, form a phthalimide ring; a phthalimide ring substituted with a C₁₋₆ alkyl group; an indole ring; an indane ring; an indazole ring; a benzotriazole ring; an S,S-dioxobenzothiophene ring; a

2,3-dihydroimidazo[2,1-b]benzothiazole ring; a dibenzofuran ring; a dibenzofuran ring substituted with a C₁₋₆ alkoxy group; a fluorene ring; a fluorene ring substituted with a halogen atom; a pyrene ring; a carbostyryl ring; a carbostyryl ring substituted with a C₁₋₆ alkyl group; a naphthalene ring; a naphthalene ring substituted with 1 to 3 substituents selected from the group consisting of cyano groups, halogen atoms, nitro groups, and C₁₋₆ alkyl groups; a 1,2,3,4-tetrahydronaphthalene ring; a quinoline ring; a quinoline ring substituted with a C₁₋₆ alkyl group; an isoquinoline ring; a 2-oxo- α -chromene ring; a 2-oxo- α -chromene ring substituted with 1 to 3 substituents selected from the group

[illegible]

2. The inhibitor for production of 20-hydroxyeicosatetraenoic acid, comprising, as an effective ingredient, a hydroxyformamidine derivative, according to Claim 1, wherein R¹ to R⁵ are identical or different and represent a hydrogen atom; a hydroxyl group; a carboxyl group; a halogen atom; a C₁₋₁₄ alkyl group; a C₁₋₁₄ alkyl group substituted with 1 to 6 halogen atoms; a C₂₋₆ alkynyl group; a C₃₋₈ cycloalkyl group; a C₃₋₈ cycloalkoxy group; a C₂₋₁₀ alkanoyl group; a C₁₋₆ hydroxyalkyl group; a C₁₋₆ hydroxyalkyl group substituted with 1 to 6 halogen atoms; a C₂₋₆ alkoxy carbonyl group; a 3-phenyl-2-propenyloxy carbonyl group; a C₂₋₆ alkoxy carbonyl C₁₋₆ alkyl group; a di(C₁₋₆ alkyl)amino C₂₋₆ alkoxy carbonyl group; a mono- or di(C₁₋₆ alkyl)amino group; a C₂₋₁₀ alkanoylamino group; a C₂₋₆ alkanoylamino group substituted with a C₁₋₆ alkyl group; a benzoylamino group; a carbamoyl group; a carbamoyl group mono- or di-substituted with C₁₋₆ alkyl or phenyl groups; an N-(N',N'-di(C₁₋₆ alkyl)amino C₁₋₆ alkyl)carbamoyl group; a cyano group; a cyano C₁₋₆ alkyl group; a nitro group; a thiol group; a phenoxy group; a phenoxy group substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, and halogen atoms; a phenylthio group; a nitrophenylthio group; a C₁₋₆ alkylsulfonyl group; a phenylsulfonyl group; a C₁₋₆ alkylthio C₁₋₆ alkyl group; a phenylsulfonyl C₁₋₆ alkylthio group wherein the benzene ring is substituted with 1 to 5 halogen atoms; a phenyl group; a benzyl group; a phenyl group substituted with 1 to 3 substituents selected from the group consisting of cyano groups, halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a biphenyl group; an α -cyanobenzyl group; an α -cyanobenzyl group

substituted with 1 to 5 halogen atoms; a benzyl group substituted with a bicyclo[2.2.1]-hept-5-en-2,3-dicarboxyimidyl group; a benzoyl group; a styryl group; a styryl group substituted with 1 to 5 substituents selected from the group consisting of C₁₋₆ alkoxy groups and di(C₁₋₆ alkyl) amino alkyl groups; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyrimidinyl group; a pyrimidinyl group substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups and C₁₋₆ alkoxy groups; a phthalimidoyl group; a phthalimidoyl group substituted with 1 to 3 halogen atoms; an N-carbazolyl group; a dioxopiperidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups; a phenylsulfonylamino group; a phenylsulfonylamino group substituted with 1 to 3 C₁₋₆ alkyl groups; a C₁₋₆ alkylaminosulfonyl C₁₋₆ alkyl group; a thiadiazolyl group; an oxadiazolyl group; an oxadiazolyl group substituted with a substituted phenyl group wherein the substituents in the substituted phenyl group are 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a pyrrolidinyl group; a pyrazolyl group; a pyrazolyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and trifluoromethyl groups; a furyl group; a furyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₂₋₆ alkoxy carbonyl groups; a thienopyrimidinylthio group; a thienopyrimidinylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a thienopyridylthio group; a thienopyridylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a benzothiazolylthio group; a benzothiazolylthio group substituted with 1 to 3 halogen atoms; or a group represented by the formula: -Y-(CR⁶¹R⁶²)_m-(CR⁶³R⁶⁴)_n-R⁷ [wherein Y represents an oxygen or sulfur atom; R⁶¹, R⁶², R⁶³, and R⁶⁴ are identical or different and represent a hydrogen atom, a halogen atom, a C₁₋₄ alkyl group, or a trifluoromethyl group; R⁷ represents a hydrogen atom; a halogen atom; a C₁₋₁₄ alkyl group; a C₃₋₈ cycloalkyl group; a C₂₋₁₀ alkenyl group; a C₂₋₆ alkynyl group;


a phenyl group; a phenyl group substituted with 1 to 3 substituents selected from the group consisting of nitro groups, cyano groups, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, phenyl groups, phenoxy groups, phenethyl groups, C₂₋₆ alkoxy carbonyl groups, and halogen atoms; a cyano group; a carboxyl group; a C₁₋₆ alkoxy group; a C₁₋₆ hydroxyalkyl group; a C₃₋₈ cycloalkoxy group; a C₁₋₆ alkoxy C₁₋₆ alkoxy group; a C₁₋₆ alkoxy C₁₋₆ alkoxy C₁₋₆ alkoxy group; a C₁₋₆ alkylthio group; a C₂₋₆ alkanoyloxy group; a C₂₋₆ alkanoyloxy C₁₋₆ alkyl group; a phenoxy group; a phenylthio group; an N-(C₁₋₆ alkyl)toluidino group; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyridyl group substituted with a C₁₋₆ alkyl group; a piperidino group substituted with a C₁₋₆ alkyl group; a pyridyl group substituted with a C₁₋₆ alkoxy group; a pyrrolidino group substituted with a C₁₋₆ alkyl group; a morpholino group substituted with a C₁₋₆ alkyl group; a morpholinyl group; a morpholinyl group substituted with a C₁₋₆ alkyl group; a homomorpholinyl group; a thiomorpholino group; a thiomorpholino group substituted with a C₁₋₆ alkyl group; a thiomorpholinyl group; a thiomorpholinyl group substituted with a C₁₋₆ alkyl group; a piperadinyl group; a piperadin-1-yl group substituted with a C₁₋₆ alkyl group at the 4-position; a homopiperidinyl group; a homopiperidinyl group substituted with a C₁₋₆ alkyl group; a pyridylthio group; a quinolyl group; a furyl group; an oxetanyl group; an oxolanyl group; a dioxolanyl group; a dioxolanyl group substituted with a C₁₋₆ alkyl group; an oxanyl group; a dioxanyl group; a dioxanyl group substituted with a C₁₋₆ alkyl group; a benzodioxanyl group; a pyrrolidon-1-yl group; a pyrrolidinyl group; an N-(C₁₋₆ alkyl)pyrrolidinyl group; a piperidinyl group; an N-(C₁₋₆ alkyl)piperidinyl group; a pyrrolyl group; a thienyl group; a thiazolyl group; a thiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups; a 2,6-purindion-7-yl group substituted with C₁₋₆ alkyl group(s); a furfuryl group; a di(C₁₋₆ alkyl)amino group; a C₂₋₆ alkoxy carbonyl group; or a di(C₁₋₆ alkyl)amino C₁₋₆ alkoxy group; m is an integer of 1 to 6; and n is an integer of 0 to 6], or a

pharmaceutically-acceptable salt thereof.

3. The inhibitor for production of 20-hydroxyeicosatetraenoic acid, comprising, as an effective ingredient, a hydroxyformamidine derivative or a pharmaceutically-acceptable salt thereof, according to Claim 2, wherein R^1 , R^2 , R^4 , and R^5 represent hydrogen atoms.

4. The inhibitor for production of 20-hydroxyeicosatetraenoic acid, according to any one of Claims 1 to 3, which is a therapeutic agent for kidney diseases, cerebrovascular diseases, or circulatory diseases.

5. A hydroxyformamidine derivative represented by the formula:



wherein at least one of R^{11} to R^{55} represents a C_{5-14} alkyl group; a C_{2-6} alkenyl group; a C_{3-8} cycloalkyl C_{1-6} alkyl group; a C_{2-6} alkynyl group; a C_{3-8} cycloalkyl group; a C_{3-8} cycloalkoxy group; a C_{2-10} alkanoyl group; a C_{1-6} hydroxyalkyl group; a C_{1-6} hydroxyalkyl group substituted with 1 to 6 halogen atoms; a C_{2-6} alkoxycarbonyl group; a 3-phenyl-2-propenyloxycarbonyl group; a C_{2-6} alkoxycarbonyl C_{1-6} alkyl group; a di(C_{1-6} alkyl)amino C_{2-6} alkoxycarbonyl group; a mono- or di(C_{1-6} alkyl)amino group; a C_{2-10} alkanoylamino group; a C_{2-6} alkanoylamino group substituted with a C_{1-6} alkyl group; a benzoylamino group; a carbamoyl group; a carbamoyl group mono- or di-substituted with C_{1-6} alkyl or phenyl groups; an N-(N' , N' -di(C_{1-6} alkyl)amino C_{1-6} alkyl) carbamoyl group; a cyano group; a cyano C_{1-6} alkyl group; a C_{1-6} alkylsulfonyl group; a phenylsulfonyl group; a C_{1-6} alkylthio C_{1-6} alkyl group; a phenylsulfonyl C_{1-6} alkylthio group wherein the benzene ring is

substituted with 1 to 5 halogen atoms; a phenyl group; a benzyl group; a phenyl group substituted with 1 to 3 substituents selected from the group consisting of cyano groups, halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a biphenyl group; an α-cyanobenzyl group; an α-cyanobenzyl group substituted with 1 to 5 halogen atoms; a benzyl group substituted with a bicyclo[2.2.1]-hept-5-en-2,3-dicarboxyimidyl group; a benzoyl group; a styryl group; a styryl group substituted with 1 to 5 substituents selected from the group consisting of C₁₋₆ alkoxy groups and di(C₁₋₆ alkyl)amino alkyl groups; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyrimidinyl group; a pyrimidinyl group substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups and C₁₋₆ alkoxy groups; a phthalimidoyl group; a phthalimidoyl group substituted with 1 to 3 halogen atoms; an N-carbazolyl group; a dioxopiperidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups; a phenylsulfonylamino group; a phenylsulfonylamino group substituted with 1 to 3 C₁₋₆ alkyl groups; a C₁₋₆ alkylaminosulfonyl C₁₋₆ alkyl group; a thiadiazolyl group; an oxadiazolyl group; an oxadiazolyl group substituted with a substituted phenyl group wherein the substituents in the substituted phenyl group are 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a pyrrolidinyl group; a pyrazolyl group; a pyrazolyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and trifluoromethyl groups; a furyl group; a furyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₂₋₆ alkoxy carbonyl groups; a thienopyrimidinylthio group; a thienopyrimidinylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a thienopyridylthio group; a thienopyridylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a benzothiazolylthio group; a benzothiazolylthio group substituted with 1 to 3 halogen atoms; a group represented by the formula: -Y-(CR⁶¹R⁶²)_m-(CR⁶³R⁶⁴)_n-R⁷⁷ [wherein Y represents an oxygen or sulfur

Sub
A9
cont

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	

Sub
A9
cont

alkyl groups; a 2,6-purindion-7-yl group substituted with at least one C₁₋₆ alkyl group; a furfuryl group; a di(C₁₋₆ alkyl) amino group; a C₂₋₆ alkoxy carbonyl group; or a di(C₁₋₆ alkyl) amino C₁₋₆ alkoxy group; m is an integer of 1 to 6; and n is an integer of 0 to 6]; or a group represented by the formula: -SO₂NR⁸R⁹ [wherein R⁸ and R⁹ are identical or different and represent a hydrogen atom, a C₁₋₁₀ alkyl group, a C₂₋₆ alkanoyl group, an isoxazolyl group, an isoxazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiadiazolyl group, a thiadiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiazolyl group, a thiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyridyl group, a pyridyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyrimidinyl group, a pyrimidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyrimidinyl group substituted with 1 to 3 C₁₋₆ alkoxy groups, a pyridazinyl group, a pyridazinyl group substituted with 1 to 3 C₁₋₆ alkoxy groups, an indazolyl group, or a carbamoyl group mono- or di-substituted with C₁₋₆ alkyl groups, or alternatively, taken together with the nitrogen atom to which they are bonded, form a 3,5-dioxopiperadino group, a pyrrolidinyl group, a piperidino group, or a morpholino group], or alternatively,

the two groups adjacent to each other of R¹¹ to R⁵⁵, taken together with the benzene ring to which they are bonded, form a phthalimide ring; a phthalimide ring substituted with a C₁₋₆ alkyl group; an indole ring; an indane ring; an indazole ring; a benzotriazole ring; an S,S-dioxobenzothiophene ring; a 2,3-dihydroimidazo[2,1-b]benzothiazole ring; a dibenzofuran ring; a dibenzofuran ring substituted with a C₁₋₆ alkoxy group; a fluorene ring; a fluorene ring substituted with a halogen atom; a pyrene ring; a carbostyryl ring; a carbostyryl ring substituted with a C₁₋₆ alkyl group; a naphthalene ring; a naphthalene ring substituted with 1 to 3 substituents selected from the group consisting of cyano groups, halogen atoms, nitro groups, and C₁₋₆ alkyl groups; a 1,2,3,4-tetrahydronaphthalene ring; a quinoline ring; a quinoline ring substituted with a C₁₋₆ alkyl group; an

Sub
A9
Cont

isoquinoline ring; a 2-oxo- α -chromene ring; a 2-oxo- α -chromene ring substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, and C₁₋₆ alkoxy C₁₋₆ alkyl groups; a cinnolin ring; a cinnolin ring substituted with a C₁₋₆ alkyl group; a phthalazindione ring; a benzothiazol ring; a benzothiazol ring substituted with a C₁₋₆ alkyl group; a benzodioxorane ring; and a benzobutyrolactone ring, and the remaining groups of R¹¹ to R⁵⁵ are identical or different and represent a hydrogen atom, a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, a trifluoromethyl group, a nitro group, or a halogen atom, or a pharmaceutically-acceptable salt thereof.

Sub
B3

6. The hydroxyformamidine derivative or a pharmaceutically-acceptable salt thereof, according to Claim 5, wherein at least one of R¹¹ to R⁵⁵ represents a C₅₋₁₄ alkyl group; a C₃₋₈ cycloalkoxy group; a C₂₋₆ alkynyl group; a C₃₋₈ cycloalkyl group; a C₂₋₁₀ alkanoyl group; a C₁₋₆ hydroxyalkyl group; a C₁₋₆ hydroxyalkyl group substituted with 1 to 6 halogen atoms; a C₂₋₆ alkoxy carbonyl group; a 3-phenyl-2-propenyloxycarbonyl group; a C₂₋₆ alkoxy carbonyl C₁₋₆ alkyl group; a di(C₁₋₆ alkyl)amino C₂₋₆ alkoxy carbonyl group; a mono- or di(C₁₋₆ alkyl)amino group; a C₂₋₁₀ alkanoylamino group; a C₂₋₆ alkanoylamino group substituted with a C₁₋₆ alkyl group; a benzoylamino group; a carbamoyl group; a carbamoyl group mono- or di-substituted with C₁₋₆ alkyl or phenyl groups; an N-(N', N'-di(C₁₋₆ alkyl)amino C₁₋₆ alkyl) carbamoyl group; a cyano group; a cyano C₁₋₆ alkyl group; a C₁₋₆ alkylsulfonyl group; a phenylsulfonyl group; a C₁₋₆ alkylthio C₁₋₆ alkyl group; a phenylsulfonyl C₁₋₆ alkylthio group wherein the benzene ring is substituted with 1 to 5 halogen atoms; a phenyl group; a benzyl group; a phenyl group substituted with 1 to 3 substituents selected from the group consisting of cyano groups, halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a biphenyl group; an α -cyanobenzyl group; an α -cyanobenzyl group substituted with 1 to 5 halogen atoms; a benzyl group substituted with a

bicyclo[2.2.1]-hept-5-en-2,3-dicarboxyimidyl group; a benzoyl group; a styryl group; a styryl group substituted with 1 to 5 substituents selected from the group consisting of C₁₋₆ alkoxy groups and di(C₁₋₆ alkyl)amino alkyl groups; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyrimidinyl group; a pyrimidinyl group substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups and C₁₋₆ alkoxy groups; a phthalimidoyl group; a phthalimidoyl group substituted with 1 to 3 halogen atoms; an N-carbazolyl group; a dioxopiperidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups; a phenylsulfonylamino group; a phenylsulfonylamino group substituted with 1 to 3 C₁₋₆ alkyl groups; a C₁₋₆ alkylaminosulfonyl C₁₋₆ alkyl group; a thiadiazolyl group; an oxadiazolyl group; an oxadiazolyl group substituted with a substituted phenyl group wherein the substituents in the substituted phenyl group are 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a pyrrolidinyl group; a pyrazolyl group; a pyrazolyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and trifluoromethyl groups; a furyl group; a furyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₂₋₆ alkoxy carbonyl groups; a thienopyrimidinylthio group; a thienopyrimidinylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a thienopyridylthio group; a thienopyridylthio group substituted with 1 to 3 C₁₋₆ alkyl groups; a benzothiazolylthio group; a benzothiazolylthio group substituted with 1 to 3 halogen atoms; or a group represented by the formula: -SO₂NR⁸R⁹ [wherein R⁸ and R⁹ are identical or different and represent a hydrogen atom, a C₁₋₁₀ alkyl group, a C₂₋₆ alkanoyl group, an isoxazolyl group, an isoxazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiadiazolyl group, a thiadiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiazolyl group, a thiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyridyl group, a pyridyl group substituted with 1 to 3 C₁₋₆ alkyl groups,

Sub
B3

7. The hydroxyformamidine derivative or a

pharmaceutically-acceptable salt thereof, according to Claim 6, wherein at least one of R¹¹ to R⁵⁵ represents a C₅₋₁₄ alkyl group; a C₂₋₆ alkynyl group; a C₃₋₈ cycloalkyl group; a C₃₋₈ cycloalkoxy group; a C₂₋₁₀ alkanoyl group; a C₁₋₆ hydroxyalkyl group; a C₁₋₆ hydroxyalkyl group substituted with 1 to 6 halogen atoms; a C₂₋₆ alkoxy carbonyl group; a 3-phenyl-2-propenyloxy carbonyl group; a C₂₋₆ alkoxy carbonyl C₁₋₆ alkyl group; a di(C₁₋₆ alkyl)amino C₂₋₆ alkoxy carbonyl group; a mono- or di(C₁₋₆ alkyl)amino group; a C₂₋₁₀ alkanoylamino group; a C₂₋₆ alkanoylamino group substituted with a C₁₋₆ alkyl group; a carbamoyl group; a carbamoyl group mono- or di-substituted with C₁₋₆ alkyl or phenyl groups; an N-(N', N'-di(C₁₋₆ alkyl)amino C₁₋₆ alkyl)carbamoyl group; a cyano group; a cyano C₁₋₆ alkyl group; a C₁₋₆ alkylsulfonyl group; a phenylsulfonyl group; a C₁₋₆ alkylthio C₁₋₆ alkyl group; a phenyl group; a benzyl group; a phenyl group substituted with 1 to 3 substituents selected from the group consisting of cyano groups, halogen atoms, C₁₋₆ alkyl groups, and C₁₋₆ alkoxy groups; a biphenyl group; an α-cyanobenzyl group; an α-cyanobenzyl group substituted with 1 to 5 halogen atoms; a benzoyl group; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyrimidinyl group; a pyrimidinyl group substituted with 1 to 3 substituents selected from the group consisting of C₁₋₆ alkyl groups and C₁₋₆ alkoxy groups; a pyrrolidinyl group; a pyrazolyl group; a pyrazolyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and trifluoromethyl groups; a furyl group; a furyl group substituted with 1 to 3 substituents selected from the group consisting of halogen atoms, C₁₋₆ alkyl groups, and C₂₋₆ alkoxy carbonyl groups; or a group represented by the formula: -SO₂NR⁸R⁹ [wherein R⁸ and R⁹ are identical or different and represent a hydrogen atom, a C₁₋₁₀ alkyl group, a C₂₋₆ alkanoyl group, an isoxazolyl group, an isoxazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiadiazolyl group, a thiadiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a thiazolyl group, a thiazolyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyridyl

Sub B3
 group, a pyridyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyrimidinyl group, a pyrimidinyl group substituted with 1 to 3 C₁₋₆ alkyl groups, a pyrimidinyl group substituted with 1 to 3 C₁₋₆ alkoxy groups, a pyridazinyl group, a pyridazinyl group substituted with 1 to 3 C₁₋₆ alkoxy groups, an indazolyl group, or a carbamoyl group mono- or di-substituted with C₁₋₆ alkyl groups, or alternatively, taken together with the nitrogen atom to which they are bonded, form a 3,5-dioxopiperadino group, a pyrrolidinyl group, a piperidino group, or a morpholino group] and the remaining groups of R¹¹ to R⁵⁵ are identical or different and represent a hydrogen atom, a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, a trifluoromethyl group, a nitro group, or a halogen atom.

8. The hydroxyformamidine derivative or a pharmaceutically-acceptable salt thereof, according to Claim 5, wherein at least one of R¹¹ to R⁵⁵ represents a group represented by the formula: $-Y-(CR^{61}R^{62})_m-(CR^{63}R^{64})_n-R^{77}$ [wherein Y represents an oxygen or sulfur atom; R⁶¹, R⁶², R⁶³, and R⁶⁴ are identical or different and represent a hydrogen atom, a halogen atom, a C₁₋₄ alkyl group, or a trifluoromethyl group; R⁷⁷ represents a halogen atom; a C₄₋₁₄ alkyl group; a C₃₋₈ cycloalkyl group; a C₂₋₁₀ alkenyl group; a C₂₋₆ alkynyl group; a phenyl group; a phenyl group substituted with 1 to 3 substituents selected from the group consisting of nitro groups, cyano groups, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylthio groups, phenyl groups, phenoxy groups, phenethyl groups, C₂₋₆ alkoxy carbonyl groups, and halogen atoms; a cyano group; a carboxyl group; a C₁₋₆ alkoxy group; a C₁₋₆ hydroxyalkyl group; a C₃₋₈ cycloalkoxy group; a C₁₋₆ alkoxy C₁₋₆ alkoxy group; a C₁₋₆ alkoxy C₁₋₆ alkoxy C₁₋₆ alkoxy group; a C₁₋₆ alkylthio group; a C₂₋₆ alkanoyloxy group; a C₂₋₆ alkanoyloxy C₁₋₆ alkyl group; a phenoxy group; a phenylthio group; an N-(C₁₋₆ alkyl) toluidino group; a pyrrolidino group; a piperidino group; a morpholino group; a pyridyl group; a pyridyl group substituted with a C₁₋₆ alkyl group; a piperidino group substituted with a C₁₋₆ alkyl group; a pyridyl group substituted

[illegible]

9. The hydroxyformamidine derivative or a pharmaceutically-acceptable salt thereof, according to Claim 8, wherein at least one of R^{11} to R^{55} represents a group represented by the formula: $-O-(CR^{61}R^{62})_m-(CR^{63}R^{64})_n-R^{77}$ [wherein R^{61} , R^{62} , R^{63} , and R^{64} are identical or different and represent a hydrogen atom, a halogen atom, a C_{1-4} alkyl group, or a trifluoromethyl group; R^{77} represents a di(C_{1-6} alkyl)amino group; a di(C_{1-6} alkyl)amino C_{1-6} alkoxy group; a piperidyl group; a piperidinyl group substituted

with a C₁₋₆ alkyl group; a piperidino group; a piperidino group substituted with a C₁₋₆ alkyl group; a pyridyl group; a pyridinyl group substituted with a C₁₋₆ alkyl group; a pyridinyl group substituted with a C₁₋₆ alkoxy group; a pyridylthio group; a pyrrolidino group; a pyrrolidino group substituted with a C₁₋₆ alkyl group; a pyrrolidon-1-yl group; a pyrrolidinyl group; a pyrrolidinyl group substituted with a C₁₋₆ alkyl group; a pyrrolyl group; a thienyl group; a thiazolyl group; a morpholino group; a morpholino group substituted with a C₁₋₆ alkyl group; a morpholinyl group; a morpholinyl group substituted with a C₁₋₆ alkyl group; a homomorpholinyl group; a thiomorpholino group; a thiomorpholino group substituted with a C₁₋₆ alkyl group; a thiomorpholinyl group; a thiomorpholinyl group substituted with a C₁₋₆ alkyl group; a piperadiny group; a piperadin-1-yl group substituted with a C₁₋₆ alkyl group at the 4-position; a homopiperidinyl group; or a homopiperidinyl group substituted with a C₁₋₆ alkyl group; m is an integer of 1 to 6; and n is an integer of 0 to 6], and the remaining groups of R¹¹ to R⁵⁵ are identical or different and represent a hydrogen atom, a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, a trifluoromethyl group, a nitro group, or a halogen atom.

10. The hydroxyformamide derivative or a pharmaceutically-acceptable salt thereof, according to any one of Claims 7 to 9, wherein R¹¹, R²², R⁴⁴, and R⁵⁵ represent hydrogen atoms.

11. An inhibitor for production of 20-hydroxyeicosatetraenoic acid, comprising, as an effective ingredient, the hydroxyformamide derivative or a pharmaceutically-acceptable salt thereof according to any one of Claims 5 to 10.

12. The inhibitor for production of 20-hydroxyeicosatetraenoic acid, according to Claim 11, which is

a therapeutic agent for kidney diseases, cerebrovascular diseases, or circulatory diseases.

add
B5

add
 p_1

[illegible]